

RATIONALITY CRITERIA FOR MOTIVIC ZETA-FUNCTIONS

MICHAEL LARSEN AND VALERY A. LUNTS

1. INTRODUCTION

Let X be a variety over a finite field $\mathbb{F}$ and $\mathrm{Sym}^n X = X^n/S_n$, the variety of effective zero-cycles on X of degree n (where by convention, $S^0 X = \mathrm{Spec} \mathbb{F}$). A celebrated theorem of B. Dwork asserts that the zeta-function

$$Z_X(t) = \sum_{i=0}^{\infty} |\mathrm{Sym}^n(X)(\mathbb{F})| t^n$$

is a rational function in t . Kapranov asked [Ka] whether this rationality lifts to the Grothendieck ring of varieties over $\mathbb{F}$. Explicitly, let $K_0[\mathcal{V}_{\mathbb{F}}]$ denote the ring of $\mathbb{Z}$ -combinations of isomorphism classes of $\mathbb{F}$ -varieties modulo the cutting-and-pasting relation

$$[X] = [Y] + [X \setminus Y]$$

for closed $\mathbb{F}$ -subvarieties $Y \subset X$. Is the **motivic zeta-function**

$$\zeta_X(t) = \sum_{i=0}^{\infty} [\mathrm{Sym}^n(X)] t^n$$

always rational as a power series in $K_0[\mathcal{V}_{\mathbb{F}}]$? More generally, is this true for varieties over a general field K , for example, $K = \mathbb{C}$? If so, this would give a fundamentally new proof of Dwork's theorem, one that does not depend on Frobenius at all. Kapranov observed that it *is* true when X is a curve. (See Theorem 3.7 below for a statement and proof over $\mathbb{C}$.)

In [LaLu], we proved that $\zeta_X(t)$ is *not* rational when X is a complex projective non-singular surface with geometric genus ≥ 2 . In this paper we return to the problem and give a simple necessary and sufficient condition on complex surfaces for $\zeta_X(t)$ to be rational:

Theorem 1.1. *A complex surface X has rational motivic zeta function if and only if it has Kodaira dimension $-\infty$.*

The notion of rationality needs some explanation (see §2 below). Theorem 3.9 asserts rationality in the strong sense when Kodaira dimension is $-\infty$, and Theorem 7.6 denies rationality in the weak sense when Kodaira

The first named author was partially supported by NSF grant DMS-0100537. The second named author was partially supported by NSA grant MDA904-01-1-0020 and CRDF grant RM1-2405-MO-02.

dimension is ≥ 0 . The methods for proving rationality and irrationality are entirely different. The fact that they meet in the middle to give a necessary and sufficient condition appears as a minor miracle, made possible by the classification of surfaces. By contrast, in dimension ≥ 3 , the rationality problem seems wide open.

There are two main ideas in this paper. The first is to probe rationality by means of new **motivic measures**, i.e. field-valued points of $\mathrm{Spec} K_0[\mathcal{V}_{\mathbb{C}}]$. In [LaLu], we showed that any multiplicative function on the set of stable birational equivalence classes of non-singular complex projective varieties defines a motivic measure. In particular, we constructed a measure μ_1 , with values in the fraction field of the group ring of the multiplicative group of integer power series $(1 + s\mathbb{Z}[[s]])^\times$, characterized by the formula

$$\mu_1([Z]) = \left[\sum_{i=0}^{\infty} \dim \Gamma(Z, \Omega_Z^i) s^i \right]$$

for Z a (connected) non-singular projective variety. In the current paper, we introduce a set of measures μ_n indexed by positive integers n , characterized by

$$(1.1) \quad \mu_n([Z]) = \left[\sum_{i=0}^{\infty} \dim H^0(Z, \Psi^n \Omega_Z^i) s^i \right],$$

where n denotes the n th Adams operation. The point of this generalization is that it allows us to work with higher plurigena in much the way that we worked with geometric genus in [LaLu]. If for some n , $\mu_n(\zeta_X(t))$ is irrational, then of course $\zeta_X(t)$ itself is irrational. For singular Z , the right hand side of (1.1) does not make sense, and indeed the left hand side is generally not of the form $[P]$ for any power series $P \in (1 + s\mathbb{Z}[[s]])^\times$. However, in the special case $Z = \mathrm{Sym}^n X$, $\dim X = 2$, it turns out that $\mu_n([Z]) = \mu_n([\mathrm{Hilb}^n X])$, where $\mathrm{Hilb}^n X$ is the Hilbert scheme of points on X . (The fact that $\mathrm{Hilb}^n X$ is non-singular is another way in which dimension 2 is special.) Our task, therefore, is to prove that if the m th plurigenus of X is positive, then

$$\sum_{i=0}^{\infty} \left[\sum_{j=0}^{\infty} \dim H^0(\mathrm{Hilb}^i X, \Psi^m \Omega_{\mathrm{Hilb}^i X}^j) s^j \right] t^i$$

is irrational.

The other main idea in this paper is to make systematic use of λ -ring ideas and techniques (see §4 for a brief review of this theory). We have already noted the appearance of Adams operations. This is somewhat delicate: we cannot work in the usual $K(X)$ since the global section functor is not well-defined there. Instead, we need to prove that the ring of virtual vector bundles modulo *split* exact sequences is a special λ -ring (see §5). Also significant is the idea that the motivic zeta-function (or better, its inverse) should be regarded as the universal λ -homomorphism. This is true only formally since $K_0[\mathcal{V}_{\mathbb{C}}]$ is not a special λ -ring. It is nonetheless suggestive,

since the universal λ -homomorphism sends every virtually finite element to a rational power series. For a special λ -ring the virtually finite elements form a λ -subring which in many interesting cases is the whole ring. This interpretation of the zeta-function suggests on the one hand that we should seek motivic measures which are λ -homomorphisms and on the other that it may be natural to replace the Grothendieck ring of varieties by its *specialization*, i.e., its maximal quotient which is a special λ -ring (see §8).

E. Looijenga has called our attention to certain formal analogies between the problem of rationality of zeta-functions of complex surfaces X , and Severi's conjecture, disproved by D. Mumford [Mu], which predicted that the group of 0-cycles modulo rational equivalence on X would be finite-dimensional. Each statement can be understood as bounding the complexity of symmetric powers of X . In each case, global holomorphic 2-forms provide the obstruction to this boundedness. However, there is an essential difference. A conjecture of S. Bloch [Bl] asserts that when $H^0(X, \Omega_X^2) = 0$, the group of degree zero 0-cycles should be isomorphic to the Albanese variety of X . This has been settled by Bloch, A. Kas, and D. Lieberman [BKL] for Kodaira dimension < 2 , so there are examples of surfaces in which the two notions of boundedness diverge. In particular, the use of plurigenera in this paper does not seem to have a counterpart in the world of CH^2 .

Throughout this paper, a **variety** will be a reduced separated scheme of finite type over $\mathbb{C}$. The class $[X] \in K_0[\mathcal{V}_{\mathbb{C}}]$ of a variety X will sometimes be written without brackets when it seems unlikely to lead to any confusion.

2. RATIONALITY CRITERIA FOR POWER SERIES

In this paper we will be concerned with the rationality of power series with coefficients in a commutative ring A . It is not entirely clear how such rationality should be defined when A is not an integral domain (and we know [Po] that the Grothendieck ring of varieties has zero-divisors). We consider several possible definitions:

Definition 2.1. A power series $f(t) \in A[[t]]$ is **globally rational** if and only if there exist polynomials $g(t), h(t) \in A[t]$ such that $f(t)$ is the unique solution of $g(t)x = h(t)$.

Definition 2.2. A power series $f(t) = \sum_i a_i t^i \in A[[t]]$ is **determinantally rational** if and only if there exist integers m and n such that

$$\det \begin{pmatrix} a_i & a_{i+1} & \cdots & a_{i+m} \\ a_{i+1} & a_{i+2} & \cdots & a_{i+m+1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{i+m} & a_{i+m+1} & \cdots & a_{i+2m} \end{pmatrix} = 0$$

for all $i > n$.

It is classical [E.B] that Definition 2.1 is equivalent to Definition 2.2 when A is a field. This suggests a third possible definition:

Definition 2.3. A power series $f \in A[[t]]$ is **pointwise rational** if and only if for all homomorphisms Φ from A to a field, $\Phi(f)$ is rational by either of the two previous definitions.

These definitions are related by the following proposition:

Proposition 2.4. *Any globally rational power series is determinantly rational, and any determinantly rational power series is pointwise rational. Neither converse holds for a general coefficient ring A . All three conditions are equivalent when A is an integral domain.*

Proof. Suppose $g(t) = \sum_i^k b_i t^i, h(t) = \sum_i^\ell c_i t^i \in A[[t]]$ and $f(t) = \sum_i a_i t^i$ is the unique solution to $g(t)x = h(t)$. Since $f(t) + a$ is not a solution when $a \neq 0$ is a constant, the annihilator of the ideal $(b_0, b_1, \dots, b_k)$ of coefficients of g is (0) . Setting $m = k, n = \ell$ in Definition 2.2,

$$\begin{pmatrix} a_i & a_{i+1} & \cdots & a_{i+m} \\ a_{i+1} & a_{i+2} & \cdots & a_{i+m+1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{i+m} & a_{i+m+1} & \cdots & a_{i+2m} \end{pmatrix} \begin{pmatrix} b_k \\ b_{k-1} \\ \vdots \\ b_0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

for $i > \ell - k$. Left multiplying both sides by the matrix of cofactors, we conclude that the determinant of the above matrix annihilates the b -column vector, which means that it is 0.

To see that the converse does not hold, consider $A = \mathbb{Z}[x]/(x^2)$. As A is countable, the set of globally rational power series over A is countable. However, any power series of the form $f(t) = xg(t)$ satisfies Definition 2.2 when $m = 1$.

Suppose $f(t) = \sum_i a_i t^i$ satisfies Definition 2.2. The same is then true for $\Phi(f)$, which is a power series over a field. For fields, however, the determinantal condition implies rationality.

To see that the converse does not hold, consider

$$A = \mathbb{Z}[x_1, x_2, \dots]/(x_1^2, x_2^2, \dots)$$

in infinitely many variables x_i . Every homomorphism from A to a field factors through the augmentation. Therefore, $f(t) = \sum_{i=1}^\infty x_i t^i$ is pointwise rational. However, the determinant

$$\det \begin{pmatrix} a_i & a_{i+1} & \cdots & a_{i+m} \\ a_{i+1} & a_{i+2} & \cdots & a_{i+m+1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{i+m} & a_{i+m+1} & \cdots & a_{i+2m} \end{pmatrix}$$

is never zero since it has a non-trivial $x_i x_{i+2} x_{i+4} \cdots x_{i+2m}$ coefficient.

Finally, when A is an integral domain, we let Φ denote the inclusion map from A to its fraction field F . If $f(t)$ is pointwise rational, it must be

rational as a power series in $F[[t]]$. Therefore, $g(t)f(t) = h(t)$ for polynomials $g(t), h(t) \in F[t]$, $g(t) \neq 0$. Clearing denominators, we may assume $g(t), h(t) \in A[t]$. As $A[t]$ is an integral domain, $f(t)$ is globally rational. $\square$

The following lemma will be useful in §3:

Lemma 2.5. *If $f(t) \in 1 + tA[[t]]$ is globally (resp. pointwise) rational, the same is true of $f(t)^{-1}$.*

Proof. If $f(t)$ is the unique solution of $g(t)x = h(t)$, then $g(t)$ is not a zero-divisor in $A[[t]]$, and $f(t)$ is invertible, so again not a zero-divisor. Therefore $h(t) = f(t)g(t)$ is not a zero divisor, and $f(t)^{-1}$ is the unique solution of $h(t)x = g(t)$. The pointwise case is trivial. $\square$

In [Ka], the rationality of motivic zeta functions is discussed in pointwise terms. To give the strongest possible results, we generally prove rationality globally and irrationality pointwise. To do the latter, we make free use of the determinantal formulation of rationality for power series over fields.

Consider a free abelian group G and its group ring $\mathbb{Z}[G]$. This ring is isomorphic to the ring of Laurent polynomials, hence is a domain. Let F be its field of fractions. Denote by Θ the collection of power series $\sum g_i t^i \in F[[t]]$, where $g_i \in G$ or $g_i = 0$. The following proposition gives a rationality criterion for power series in Θ .

Proposition 2.6. *Let $f(t) = \sum g_i t^i \in \Theta$ be a power series in $F[[t]]$. Then $f(t)$ is rational if and only if there exists $n \geq 1$, $i_0 \in \mathbb{N}$ and a sequence $\{h_i \in G\}$ periodic with period n such that for $i > i_0$ we have $g_{i+n} = h_i g_i$.*

Proof. If there exist n , i_0 , and h_i as above, then

$$f(t) = \sum_{i=0}^{i_0} g_i t^i + \sum_{i=1}^n \frac{g_{i_0+i}}{1 - h_i t^n}$$

is a rational function with denominator of degree $m \leq n^2$.

For the converse, by assumption there exists a polynomial

$$q(t) = a_m t^m + a_{m-1} t^{m-1} + \cdots + a_0 \in F[t]$$

such that $q(t)f(t)$ is a polynomial. Clearing denominators, we may assume $a_i \in \mathbb{Z}[G]$ for all i . Let $C \subset G$ be the collection $\alpha\beta^{-1}$ where α, β are group elements which have non-zero coefficient in some a_j , $0 \leq j \leq m$. Then for $i > i_0$

$$(2.1) \quad -a_0 g_{i+m} = a_1 g_{i+m-1} + \cdots + a_m g_i$$

and in particular

$$(2.2) \quad g_{i+m} \in g_i C \cup \cdots \cup g_{i+m-1} C \cup \{0\}$$

Definition 2.7. A power series $p(x) = \sum \alpha_i t^i \in \Theta$ is *compact* (or (m, K) -compact) if there exists a finite set $K \subset G$ such that for nonzero α_i, α_j such that $|i - j| \leq m$ we have $\alpha_i \alpha_j^{-1} \in K$.

Case 1. Assume that $f(t)$ is compact. Fix $i > i_0$. Using the compactness of $f(t)$ we can find $n \geq 1$ and $h \in G$ such that

$$(g_{i+n}, g_{i+n+1}, \dots, g_{i+n+m-1}) = h(g_i, g_{i+1}, \dots, g_{i+m-1}).$$

Then (2.1) implies

$$(g_{j+n}, g_{j+n+1}, \dots, g_{j+n+m-1}) = h(g_j, g_{j+1}, \dots, g_{j+m-1})$$

for all $j \geq i$, which proves the proposition in this case. Note that we can bound n by some function $\phi(m, |K|)$.

Case 2. This is the general case. It follows from (2.2) that there exists an integer r , $1 \leq r \leq m$ and compact power series $f_1(t), \dots, f_r(t)$ in Θ such that $f(t) = \sum f_j(t)$. We choose the *minimal* such r and corresponding series $f_j(t) = \sum g_i^j t^i$ for $j = 1, 2, \dots, r$.

Lemma 2.8. *The power series $f_j(t)$ simultaneously satisfy the recurrence relations (2.1) for coefficients with indices contained in arbitrarily long intervals.*

Proof. We use the minimality of r . If $r = 1$ we are in Case 1 and there is nothing to prove. Assume $r \geq 2$. The minimality of r implies (and actually is equivalent to) the following:

For every finite subset $S \subset G$ there exist infinitely many intervals $I \subset \mathbb{N}$ of length $m + 1$ such that for any nonzero $g_{i_1}^k, g_{i_2}^l$ with $i_1, i_2 \in I$ and $k \neq l$ we have $g_{i_1}^k (g_{i_2}^l)^{-1} \notin S$.

In particular, we can find such an interval $I = (i, i + 1, \dots, i + m)$ for the set $S = C$ so that $i > i_0$. Then the recurrence relation (2.1) implies the same relation for coefficients $g_i^j, \dots, g_{i+m}^j$ for all $j = 1, \dots, r$. To get a long interval J , say of length sm on which this recurrence holds for all j we proceed as follows: let $K \subset G$ be finite, such that each $f_j(t)$ is (m, K) -compact; take $S = CK^{2s}$ and let $I = (i, i + 1, \dots, i + m)$ be a corresponding interval such that $i > i_0$. Then we can take $J = (i, i + 1, \dots, i + sm)$. This proves the lemma. $\square$

We now complete the proof of the proposition as follows: Choose a sufficiently long interval $J \subset (i_0, \infty)$ as in the proof of last lemma. Then repeating the argument in Case 1 for each $j = 1, \dots, r$ we find $n_j \geq 1$ and $h'_j \in G$ such that

$$(g_{i+n_j}^j, g_{i+n_j+1}^j, \dots, g_{i+n_j+m-1}^j) = h'_j(g_i^j, g_{i+1}^j, \dots, g_{i+m-1}^j)$$

as long as $[i + n_j + m] \subset J$. Now take $n = n_1 \cdots n_r$ and put

$$h_i = \begin{cases} (h'_j)^{n/n_j}, & \text{if } i \in J \text{ and } g_i^j \neq 0, \\ 1, & \text{if } i \in J \text{ and all } g_i^j = 0. \end{cases}$$

It follows that $g_{i+n} = h_i g_i$ as long as the indices stay in the interval J .

The recurrence (2.1) implies that there exists a linear operator T such that

$$(g_{i+n}, g_{i+n+1}, \dots, g_{i+n+m}) = T^n(g_i, g_{i+1}, \dots, g_{i+m})$$

for all $i \geq i_0$. We claim that the relation $g_{i+n} = h_i g_i$ holds in fact for all $i \geq \inf(J)$. Indeed, this follows from the following simple fact in linear algebra: Let V be a d -dimensional vector space and $0 \neq v \in V$; assume that we have two linear operators A and B in V , such that $A^l v = B^l v$ for $l = 1, \dots, d-1$. Then $A^l v = B^l v$ for all $l \geq 0$. $\square$

3. RATIONALITY THEOREMS

In this section we describe several classes of varieties which have rational zeta-functions, notably rational and ruled surfaces and linear algebraic groups. These are rather special cases, and indeed we suspect that except in dimension 1, rationality is the exception rather than the rule. Some evidence for this point of view is given in the discussion of irrationality theorems in §7.

Lemma 3.1. *If X is a variety and Y a closed subvariety with complement U , then if any two of $\zeta_X(t)$, $\zeta_Y(t)$, and $\zeta_U(t)$ are globally (resp. pointwise) rational, then the third is so as well.*

Proof. Stratifying $\text{Sym}^n X$ according to how many points land in Y , we obtain

$$(3.1) \quad [\text{Sym}^n X] = \sum_{i+j=n} [\text{Sym}^i Y \times \text{Sym}^j U] = \sum_{i+j=n} [\text{Sym}^i Y][\text{Sym}^j U].$$

It follows that $\zeta_X(t) = \zeta_Y(t)\zeta_U(t)$.

By Lemma 2.5, not only are the two specified zeta-functions globally (resp. pointwise) rational, the same is true of their reciprocals. The product of rational zeta-functions is again rational, and the lemma follows. $\square$

In particular, the disjoint union of varieties with globally rational zeta-functions again has a globally rational zeta-function. This shows in particular that zero-dimensional varieties have globally rational zeta-functions. Also, a stratified variety has such a zeta-function as long as all of its strata do.

Lemma 3.2. *If X is any variety, and $E \rightarrow X$ is a vector bundle of rank r , then*

$$[E] = [X][\mathbb{A}^r].$$

Proof. Vector bundles are locally trivial in the Zariski topology, so there exists a dense open subset $U \subset E$ over which E restricts to a trivial bundle. If Y denotes the complement of U and E_U and E_Y the pull-back of E to U and Y respectively, then applying Lemma 3.1 to $E_Y \subset E$ and $Y \subset X$, it suffices to prove the proposition for Y and U , and for U , $E_U = U \times \mathbb{A}^r$. The lemma follows by Noetherian induction. $\square$

The following proposition is due to B. Totaro [Go]:

Proposition 3.3. *If X is any variety, $E \rightarrow X$ is a vector bundle of rank r , and n is any positive integer, then $[\mathrm{Sym}^n E] = [\mathrm{Sym}^n X][\mathbb{A}^1]^{rn}$. Equivalently, $\zeta_E(t) = \zeta_X(\mathbb{A}^r t)$.*

Corollary 3.4. *If X is a variety such that $\zeta_X(t)$ is globally (resp. pointwise) rational, and $E \rightarrow X$ is a vector bundle, then $\zeta_E(t)$ is globally (resp. pointwise) rational.*

Corollary 3.5. *For all non-negative integers n , $\mathbb{A}^n$ is rational.*

Corollary 3.6. *If X is a variety and $P \rightarrow X$ is a projective space bundle of rank r which is locally trivial in the Zariski topology, then*

$$\zeta_P(t) = \zeta_X(t) \zeta_X(\mathbb{A}^1 t) \cdots \zeta_X(\mathbb{A}^n t).$$

In particular, $\zeta_{\mathbb{P}^r}(t)$ is rational for all r .

Proof. By Noetherian induction, it suffices to consider the case $P = X \times \mathbb{P}^r$. The corollary then follows immediately from the stratification of $\mathbb{P}^r$ with strata $\mathbb{A}^i$, $0 \leq i \leq r$. $\square$

We come now to the main positive result in the subject.

Theorem 3.7. (Kapranov) *If X is any one-dimensional variety, then $\zeta_X(t)$ is globally rational.*

Proof. Let X be a 1-dimensional variety. The singular locus Y is either empty or zero-dimensional. In the latter case, the rationality question reduces to the case of $X \setminus Y$, so without loss of generality we may assume X is non-singular. If X has more than one component, it suffices to prove that each one has a rational zeta-function, so without loss of generality we may assume X is connected and therefore irreducible. Let $\bar{X}$ denote the unique projective non-singular curve containing X as an open subvariety. As $\bar{X} \setminus X$ is empty or zero-dimensional, without loss of generality we may assume that X is projective and non-singular.

Let g be the genus of X . Let x_0 denote a base point of X and J denote the Jacobian variety $\mathrm{Jac}^0(X)$. For non-negative $n \geq 2g - 1$, the morphism $X^n \rightarrow J$ mapping

$$(x_1, \dots, x_n) \mapsto -nx_0 + \sum_{i=1}^n x_i$$

factors through $\mathrm{Sym}^n X$ realizing it as a projective space bundle of rank $n - g$ over J . The closed immersion $X^n \rightarrow X^{n+1}$ sending $(x_1, \dots, x_n)$

to $(x_0, \dots, x_n)$ induces a closed J -immersion $\mathrm{Sym}^n X \rightarrow \mathrm{Sym}^{n+1}(X)$; the complement is a vector bundle of rank $n+1-g$ over J . By Lemma 3.2,

$$[\mathrm{Sym}^{n+1} X] - [\mathrm{Sym}^n X] = [X][\mathbb{A}^1]^{n+1-g}.$$

This implies that

$$(3.2) \quad \zeta_X(t)(1-t)(1-\mathbb{A}^1 t)$$

is a polynomial of degree $\leq 2g$. $\square$

Corollary 3.8. *If X is an algebraic surface, the global (resp. pointwise) rationality of $\zeta_X(t)$ depends only on the birational equivalence class of X .*

Proof. If X_1 and X_2 are two such surfaces and U is a surface which is isomorphic to dense open subvarieties of each, then setting $Y_i = X_i \setminus U$,

$$\zeta_{X_i}(t) = \zeta_U(t)\zeta_{Y_i}(t).$$

By Theorem 3.7, $\zeta_{Y_i}(t)$ is globally (therefore also pointwise) rational, so by Lemma 3.1, each $\zeta_{X_i}(t)$ is rational if and only if $\zeta_U(t)$ is so. $\square$

Theorem 3.9. *If X is a surface with Kodaira dimension $-\infty$, then $\zeta_X(t)$ is globally rational.*

Proof. There are two cases: rational surfaces and birationally ruled surfaces. In each case, we may choose any variety in the given birational equivalence class. For rational surfaces, we use $\mathbb{A}^2$, which has a globally rational zeta-function by Corollary 3.5. A ruled surface is a projective line bundle over a curve. By the Tsen-Lang theorem, such a $\mathbb{P}^1$ -bundle is Zariski-locally trivial, and by Theorem 3.7 and Corollary 3.6, the motivic zeta-function of a ruled surface is globally rational. $\square$

Theorem 3.10. *If X is a linear algebraic group, its zeta-function is globally rational.*

Proof. The components of an algebraic group G are isomorphic to one another as varieties, so

$$\zeta_G(t) = \zeta_{G^\circ}(t)^{[G:G^\circ]},$$

where G° denotes the identity component of G . We therefore assume that G is connected. Let U denote the unipotent radical of G and $H = G/U$. Then U is isomorphic to a subgroup of the unitriangular matrices $n \times n$ matrices [A.B] 4.8. The logarithm map therefore defines an isomorphism of varieties between U and its Lie algebra, which means that U is isomorphic to $\mathbb{A}^{\dim U}$. The existence of Levi decompositions gives an isomorphism $G \simeq U \times H$, so it suffices to prove that $\zeta_H(t)$ is globally rational.

Let B denote a Borel subgroup of H , T a maximal torus of B , and V the unipotent radical of B . Decomposition into Schubert cells gives a stratification of the flag variety H/B in which every stratum is isomorphic to $\mathbb{A}^k$ for some k with pre-image $VwB \subset H$ isomorphic to $\mathbb{A}^k \times B \cong \mathbb{A}^{k+\dim V} \times T$ [A.B] 14.12. It therefore suffices to prove that $\zeta_T(t)$ is rational.

We use induction on $d = \dim T$. The rationality is trivial for $d = 0$. Now,

$$\mathbb{G}_m^d \times \mathbb{A}^1 \setminus \mathbb{G}_m^d = \mathbb{G}_m^{d+1},$$

so by Corollary 3.4 the rationality of $\zeta_{\mathbb{G}_m^d}(t)$ implies that of $\zeta_{\mathbb{G}_m^{d+1}}(t)$. $\square$

4. LAMBDA RINGS

In this section, we develop some basic definitions and facts connected with the notion of λ -ring. A good reference for this material is the first section of [AT]

Definition 4.1. A λ -**structure** on a commutative ring A is an infinite sequence $\lambda^0, \lambda^1, \lambda^2, \dots$, of maps $A \rightarrow A$ such that

$$\begin{aligned} \lambda^0(x) &= 1 \\ \lambda^1(x) &= x \\ \lambda^n(x+y) &= \sum_{i+j=n} \lambda^i(x) \lambda^j(y). \end{aligned}$$

A λ -**ring** is a commutative ring endowed with a λ -structure. We call a ring homomorphism between λ -rings which commutes with λ -operations a λ -**homomorphism**.

The prototype of a λ -ring is the Grothendieck group of finite dimensional vector spaces over a field. Of course, explicitly the ring here is $\mathbb{Z}$. More generally, one can look at Grothendieck groups of finitely generated projective modules over a ring or finite rank vector bundles over variety or over a topological space. In each case, λ^i should be regarded as the i th exterior power operation. For example, in the first case, $\lambda^i(n) = \binom{n}{i}$.

Example 4.2. Consider the polynomial ring $\mathbb{Z}[s]$. If we identify $\mathbb{Z}$ with the ring of virtual finite dimensional vector spaces, then $\mathbb{Z}[s]$ is identified with the ring of isomorphism classes of $\mathbb{N}$ -graded virtual finite dimensional vector spaces. Using this identification define the λ -structure on $\mathbb{Z}[s]$ as follows:

$$\lambda^i(Vs^p) = \begin{cases} \text{Sym}^i V s^{ip}, & \text{if } p \text{ is even,} \\ \Lambda^i V s^{ip}, & \text{if } p \text{ is odd.} \end{cases}$$

Here Λ and Sym have their usual meanings for ordinary virtual vector spaces:

$$\begin{aligned} \Lambda^k(V - W) &= \sum_{i+j=k} (-1)^j \Lambda^i V \otimes \text{Sym}^j W; \\ \text{Sym}^k(V - W) &= \sum_{i+j=k} (-1)^j \text{Sym}^i V \otimes \Lambda^j W. \end{aligned}$$

Note that the sign conventions for λ^i are those one expects for *symmetric* powers of virtual super vector spaces.

Example 4.3. Let A be a λ -ring and $M \subset A$ a multiplicative submonoid closed under the λ -operations. Consider the corresponding monoid ring $\mathbb{Z}[M]$. (Note that it is *not* a subring of A). Then $\mathbb{Z}[M]$ has a natural λ -structure given by

$$\lambda^i([m]) = [\lambda^i m].$$

Example 4.4. Let us combine the two previous examples. Let $M \subset \mathbb{Z}[s]$ be the multiplicative monoid, which consists of polynomials with constant term 1. Then $\mathbb{Z}[M]$ is a λ -ring. This example will be important to us since our motivic measures will take their values in $\mathbb{Z}[M]$. We will also need the following lemma.

Lemma 4.5. *The monoid M is a free commutative monoid. The ring $\mathbb{Z}[M]$ is a polynomial ring, hence an integral domain.*

Proof. The ring $\mathbb{Z}[s]$ is factorial and any element of M is a unique product of elements of M , which are prime in $\mathbb{Z}[s]$ (the only unit in M is 1). Thus M is isomorphic to the monoid $\oplus \mathbb{N}$, where the summation is over all prime elements of $\mathbb{Z}[s]$ which are in M . Hence $\mathbb{Z}[M]$ is a polynomial ring, so it is an integral domain. $\square$

Example 4.6. By (3.1), the Grothendieck ring of varieties $K_0[\mathcal{V}_K]$ has a natural λ -ring structure for any field K . Indeed, put $\lambda^i[X] := [\mathrm{Sym}^i X]$. Then

$$\lambda_t([X]) = \sum_{i=0}^{\infty} \lambda^i[X] t^i = \zeta_X(t).$$

For any commutative ring A there is a natural λ -ring structure on the set $1 + tA[[t]]$. The operation of addition in this ring is given by the multiplication of power series. Multiplication $(\cdot)$ and the λ -operations (Λ^i) are given by universal polynomials which are uniquely characterized by the identities

$$(4.1) \quad \left\{ \prod_{i=1}^m (1 + a_i t) \right\} \cdot \left\{ \prod_{j=1}^n (1 + b_j t) \right\} = \prod_{i=1}^m \prod_{j=1}^n (1 + a_i b_j t)$$

and

$$\Lambda^k \prod_{i=1}^n (1 + a_i t) = \prod_{S \subset \{1, \dots, n\}, |S|=k} (1 + t \prod_{j \in S} a_j).$$

Equation (4.1) implies that the polynomial expressing the t^p coefficient of $\sum_i x_i t^i \cdot \sum_j y_j t^j$ in terms of x_i and y_j lies in the ideal

$$(x_{m+1}, x_{m+2}, \dots, y_{n+1}, y_{n+2}, \dots)$$

whenever $p > mn$. This implies that the product of two polynomials in $1 + tA[[t]]$ is again a polynomial, regardless of whether the polynomials split into linear factors. As $\cdot$ distributes over the usual multiplication of power series, $f(t) \cdot g(t)$ is a ratio of polynomials in $1 + tA[[t]]$ if $f(t)$ and $g(t)$ are.

If A has a λ -structure then the map

$$\lambda_t: A \rightarrow 1 + tA[[t]], \quad \lambda_t(a) = \sum_{i=0}^{\infty} \lambda^i a t^i$$

is an additive group homomorphism.

Definition 4.7. An element $a \in A$ is **finite dimensional** if $\lambda_t(a)$ is a polynomial, and the **dimension** of a is the degree of this polynomial. A difference of finite dimensional elements is **virtually finite**. A λ -ring is **finite dimensional** if all of its elements are virtually finite.

Definition 4.8. We say A is a **special** λ -ring if the homomorphism of additive groups λ_t is a λ -homomorphism. In this case, we will call λ the universal λ -homomorphism.

For example, the Grothendieck group of vector spaces (or projective modules or vector bundles) is a special λ -ring. Moreover, $1 + tA[[t]]$ is always special, regardless of whether or not A is so. On the other hand, Examples 4.2, 4.4, and 4.6 are not special, and neither is Example 4.3 in general, even if A happens to be so. Special λ -rings are characterized by identities of the form

$$(4.2) \quad \lambda^n(xy) = P_n(x, \lambda^2 x, \dots, \lambda^n x, y, \dots, \lambda^n y)$$

and

$$(4.3) \quad \lambda^m \lambda^n x = Q_{m,n}(x, \lambda^2 x, \dots, \lambda^{mn} x)$$

for certain universal polynomials $P_n, Q_{m,n}$. Equation (4.2) guarantees that λ_t is a ring homomorphism and (4.3) guarantees it respects λ -structures.

Lemma 4.9. *For any λ -ring A there exists a universal pair consisting of a special λ -ring B and a λ -homomorphism $A \rightarrow B$ such that every λ -homomorphism from A to a special λ -ring C factors through B .*

Proof. If C is a special λ -ring and $\Phi: A \rightarrow B$ a λ -homomorphism, then for all $x, y \in A$,

$$\begin{aligned} \Phi(\lambda^n(xy)) &= \lambda^n \Phi(xy) = \lambda^n(\Phi(x)\Phi(y)) \\ &= P_n(\Phi(x), \dots, \lambda^n \Phi(x), \Phi(y), \dots, \lambda^n \Phi(y)) \\ &= P_n(\Phi(x), \dots, \Phi(\lambda^n x), \Phi(y), \dots, \Phi(\lambda^n y)) \\ &= \Phi(P_n(x, \dots, \lambda^n x, y, \dots, \lambda^n y)), \end{aligned}$$

so

$$(4.4) \quad \lambda^n(xy) - P_n(x, \dots, \lambda^n x, y, \dots, \lambda^n y) \in \ker \Phi.$$

Similarly,

$$(4.5) \quad \lambda^m \lambda^n x - Q_{m,n}(x, \dots, \lambda^{mn} x) \in \ker \Phi.$$

Let I denote the λ -ideal in A generated by elements of type (4.4) and (4.5). Thus $I \subset \ker \Phi$. Conversely, if $B = A/I$, (4.4) and (4.5) imply (4.2) and (4.3) respectively. Thus, the quotient map $A \rightarrow B$ is universal. $\square$

We call the ring B , the **specialization** of A .

Next we say a few words about the relationship between symmetric and exterior powers. In the case that A is the Grothendieck group of vector spaces (resp. projective modules, resp. vector bundles), one-dimensional spaces (resp. invertible ideals, resp. line bundles) correspond to elements a such that $\lambda_t(a) = 1 + at$. If $b = a_1 + \cdots + a_n$, where each a_i satisfies this condition,

$$\lambda_t(b) = \prod_{i=1}^n \lambda_t(a_i) = \prod_{i=1}^n (1 + a_i t).$$

Thus,

$$\prod_{i=1}^n (1 + a_i t + a_i^2 t^2 + \cdots) = \prod_{i=1}^n \frac{1}{1 - a_i t} = \lambda_{-t}(b)^{-1}.$$

We therefore define $\sigma^n(x)$ as the n th coefficient of $\lambda_{-t}(x)^{-1}$ for all $x \in A$. We note that if

$$(4.6) \quad \sigma_t(x) = \sum_{n=0}^{\infty} \sigma^n(x) t^n = \lambda_{-t}(x)^{-1},$$

then

$$\sigma_t(x + y) = \lambda_{-t}(x + y)^{-1} = \lambda_{-t}(x)^{-1} \lambda_{-t}(y)^{-1} = \sigma_t(x) \sigma_t(y).$$

We conclude that

$$\begin{aligned} \sigma^0(x) &= 1 \\ \sigma^1(x) &= x \\ \sigma^n(x + y) &= \sum_{i+j=n} \sigma^i(x) \sigma^j(y), \end{aligned}$$

so the σ^n give a new λ -ring structure on R which we call the **opposite** structure to $\{\lambda^n\}$.

Several remarks on this construction are in order. The automorphism on the multiplicative group of power series with constant term 1 given by $f(t) \mapsto f(-t)^{-1}$ is an involution, so the opposite of the opposite of a λ -structure is the structure itself. By (4.6), an element is virtually finite with respect to a λ -structure if and only if it is virtually finite with respect to the opposite structure. The opposite of a special λ -ring need not be special. For example, in $\mathbb{Z}$, which is special with respect to $\{\lambda^n\}$, $\sigma^n(r) = \binom{r+n-1}{n}$, so the $\{\sigma^n\}$ counterpart of the identity (4.2) for $n = 2$, namely

$$\sigma^2(xy) = y^2 \sigma^2 x + x^2 \sigma^2 y - 2(\sigma^2 x)(\sigma^2 y)$$

does not hold.

The Newton polynomials, giving power sums in terms of elementary symmetric functions, allow us to define the **Adams operations** $\Psi^n: A \rightarrow A$ on any λ -ring. In fact, we need these operations only for vector bundles over algebraic varieties, where they have been extensively studied.

5. SPECIAL λ -RING $\overline{K}(X)$

Let X be a variety. It is well known that the usual K -theory (of algebraic vector bundles) associates to X a special λ -ring $K(X)$. We will need to consider a different Grothendieck group $\overline{K}(X)$ of vector bundles, so that the functor of global sections descends to a group homomorphism

$$H^0: \overline{K}(X) \rightarrow K[\text{Vect}].$$

So let us take $\overline{K}(X)$ to be the abelian group generated by isomorphism classes of algebraic vector bundles on X with relations

$$[P] = [M] + [N],$$

whenever the vector bundles $M \oplus N$ and P are isomorphic. Note that we do not impose relations coming from general short exact sequences as in the usual K -theory. The $\otimes$ operation makes $\overline{K}(X)$ a ring and the λ -operations are defined in the usual way using the exterior powers:

$$\lambda^i[P] = [\Lambda^i P].$$

Theorem 5.1. *The λ -ring $\overline{K}(X)$ is special.*

Proof. The traditional way to prove that $K(X)$ is a special λ -ring uses the splitting principle, which, in fact, is equivalent to the λ -ring (in which every element is of finite dimension) being special [FL] Ch. 1. The usual method of splitting an algebraic vector bundle produces only a short exact sequence and therefore is not applicable to our group $\overline{K}(X)$. We will prove the identities (4.2) and (4.3) in $\overline{K}(X)$ by showing that for any $x, y \in \overline{K}(X)$ there exists a λ -homomorphism from a special λ -ring to $\overline{K}(X)$ such that x, y are contained in the image. For this we need a *free* special λ -ring on two generators.

First recall the free special λ -ring on one generator. It has two standard descriptions: as a ring of symmetric functions (in an infinite number of variables), and as a direct sum of representation rings of the symmetric group. It is the second description which is useful for our purposes, so we recall it.

Let R_n be the representation ring of the symmetric group S_n , with the convention that $S_0 = \{e\}$. That is, R_n is a free $\mathbb{Z}$ -module with basis consisting of isomorphism classes of irreducible (complex) representations of S_n . Put

$$R := \bigoplus_{n \geq 0} R_n.$$

Thus R has a $\mathbb{Z}$ -basis consisting of pairs (n, ω) , where n is a natural number and ω is an irreducible representation of S_n . The ring structure is uniquely determined by the requirement that

$$(n_1, \omega_1)(n_2, \omega_2) = (n_1 + n_2, \text{Ind}_{S_{n_1} \times S_{n_2}}^{S_{n_1+n_2}} \omega_1 \otimes \omega_2).$$

The trivial $S_0 = \{e\}$ -module is the unit in the ring R . The λ -ring structure is defined as follows. Fix a basis element (n, ω) and a positive integer r . Consider the obvious action of the wreath product $S_n^r \rtimes S_r$ on $\omega^{\otimes r}$ and let $\text{Sign}(S_r)$ be the sign representation of S_r . Then $\omega^{\otimes r} \otimes \text{Sign}(S_r)$ is a left $S_n^r \rtimes S_r$ -module.

Then

$$\lambda^r(n, \omega) = (rn, \text{Ind}_{S_n^r \rtimes S_r}^{S_{rn}} \omega^{\otimes r} \otimes \text{Sign}(S_r)).$$

The special λ -ring R is generated by the element $(1, \mathbb{C})$: the elements $\{\lambda^r(1, \mathbb{C}) \mid r \geq 1\}$ are algebraically independent in R . The ring R is free in the following sense: given a special λ -ring Q and an element $x \in Q$ there exists a λ -homomorphism $f : R \rightarrow Q$ such that $f((1, \mathbb{C})) = x$.

Consider the special λ -ring $R^2 := R \otimes_{\mathbb{Z}} R$. It is the *free* special λ -ring on two generators in the obvious sense. Naturally R^2 has a $\mathbb{Z}$ -basis consisting of elements $((n_1, n_2), \omega_1 \otimes \omega_2)$, where ω_i is an irreducible representation of S_{n_i} , $i = 1, 2$; we regard $\omega_1 \otimes \omega_2$ as a $S_{n_1} \times S_{n_2}$ representation in the usual way. The λ -operations are similar to those in R :

$$\lambda^r((n_1, n_2), \omega_1 \otimes \omega_2) = ((rn_1, rn_2), \text{Ind}_{(S_{n_1}^r \times S_{n_2}^r) \rtimes S_r}^{S_{rn_1} \times S_{rn_2}} \omega_1^{\otimes r} \otimes \omega_2^{\otimes r} \otimes \text{Sign}(S_r)).$$

Let M, N be (virtual) vector bundles on X . Consider $M^{\otimes n_1} \otimes N^{\otimes n_2}$ as a (virtual) right $S_{n_1} \times S_{n_2}$ -module. We define the homomorphism $\theta : R^2 \rightarrow \overline{K}(X)$ as follows.

$$\theta((n_1, n_2), \omega_1 \otimes \omega_2) = (M^{\otimes n_1} \otimes N^{\otimes n_2}) \otimes_{S_{n_1} \times S_{n_2}} (\omega_1 \otimes \omega_2).$$

This is a ring homomorphism. We only need to check that the λ -operations correspond under θ ; that is, we need to compare

$$\begin{aligned} & \theta(\lambda^r((n_1, n_2), \omega_1 \otimes \omega_2)) \\ &= (M^{\otimes rn_1} \otimes N^{\otimes rn_2}) \otimes_{S_{rn_1} \times S_{rn_2}} \\ & \quad (\mathbb{C}[S_{rn_1} \times S_{rn_2}] \otimes_{(S_{n_1}^r \times S_{n_2}^r) \rtimes S_r} (\omega_1^{\otimes r} \otimes \omega_2^{\otimes r} \otimes \text{Sign}(S_r))) \\ &= (M^{\otimes rn_1} \otimes N^{\otimes rn_2}) \otimes_{(S_{n_1}^r \times S_{n_2}^r) \rtimes S_r} (\omega_1^{\otimes r} \otimes \omega_2^{\otimes r} \otimes \text{Sign}(S_r)) \end{aligned}$$

and

$$\begin{aligned} & \Lambda^r(\theta((n_1, n_2), \omega_1 \otimes \omega_2)) \\ &= \Lambda^r((M^{\otimes n_1} \otimes N^{\otimes n_2}) \otimes_{S_{n_1} \times S_{n_2}} (\omega_1 \otimes \omega_2)) \\ &= ((M^{\otimes rn_1} \otimes N^{\otimes rn_2}) \otimes_{S_{n_1}^r \times S_{n_2}^r} (\omega_1^{\otimes r} \otimes \omega_2^{\otimes r})) \otimes_{S_r} \text{Sign}(S_r). \end{aligned}$$

The following lemma implies the two are isomorphic.

Lemma 5.2. *Let G be a group, P, Q – right and left G -modules respectively. Consider the space $(P \otimes_G Q)^{\otimes r} = P^{\otimes r} \otimes_{G^r} Q^{\otimes r}$ as a right S_r -module. There is a canonical isomorphism of right S_r -modules*

$$(5.1) \quad \alpha: P^{\otimes r} \otimes_{G^r} Q^{\otimes r} \xrightarrow{\sim} P^{\otimes r} \otimes_{G^r \rtimes S_r} (Q^{\otimes r} \otimes \mathbb{C}[S_r]).$$

In particular there is a canonical isomorphism of vector spaces

$$(5.2) \quad \Lambda^r((P \otimes_G Q)^{\otimes r}) \xrightarrow{\sim} P^{\otimes r} \otimes_{G^r \rtimes S_r} (Q^{\otimes r} \otimes \text{Sign}(S_r)).$$

Proof of the lemma. Indeed,

$$\alpha: x \otimes y \mapsto x \otimes y \otimes 1,$$

and

$$p_1 \otimes \cdots \otimes p_r \otimes q_1 \otimes \cdots \otimes q_r \otimes \tau \mapsto p_{\tau(1)} \otimes \cdots \otimes p_{\tau(r)} \otimes q_{\tau(1)} \otimes \cdots \otimes q_{\tau(r)}$$

define mutually inverse S_r -linear maps. This proves (5.1). Tensoring both sides over S_r with $\text{Sign}(S_r)$, we get (5.2) and therefore the proposition. $\square$

So we have constructed a contravariant functor

$$\overline{K}: \{\text{Varieties}\} \rightarrow \{\text{Special } \lambda\text{-rings}\}.$$

The Adams operations Ψ^n are defined in $\overline{K}(X)$ in the usual way: these are certain universal polynomials with integer coefficients in the operations λ^i . We have

$$\Psi^n[\mathcal{L}] = [\mathcal{L}^{\otimes n}],$$

if $\mathcal{L}$ is a line bundle. Since the λ -ring $\overline{K}(X)$ is special, the Ψ^n are λ -homomorphisms (see [AtiTa], Section 5).

Note that we have a well defined group homomorphism

$$H^0: \overline{K}(X) \rightarrow K_0[\text{Vect}].$$

6. MOTIVIC MEASURES μ_n

Consider the (λ) -ring $\mathbb{Z}[M]$ as in Example 4.4. We will freely consider elements of the monoid M either as polynomials with integer coefficients or as (isomorphism classes of) graded vector spaces.

For a smooth connected projective variety X of dimension d define

$$\mu_n(X) := 1 + h_n^1(X)s + \cdots + h_n^d(X)s^d \in \mathbb{Z}[M],$$

where

$$h_n^i(X) = \dim H^0(X, \Psi^n \Omega_X^i) \in \mathbb{Z}.$$

We constructed the measure $\mu = \mu_1$ in our previous paper [LaLu]. Explicitly

$$\mu(X) = 1 + h^{1,0}(X)s + \cdots + h^{d,0}(X)s^d$$

for a smooth projective irreducible X of dimension d .

Proposition 6.1. *For each $n \geq 1$ we have*

- i) $\mu_n(X) = \mu_n(\tilde{X})$ if X and $\tilde{X}$ are birational,
- ii) $\mu_n(X \times Y) = \mu_n(X)\mu_n(Y)$,
- iii) $\mu_n(\mathbb{P}^k) = 1$ for all $k \geq 0$.

Corollary 6.2. *The mapping μ_n extends (uniquely) to a ring homomorphism*

$$\mu_n : K_0[\mathcal{V}_{\mathbb{C}}]/(\mathbb{A}^1) \rightarrow \mathbb{Z}[M].$$

Indeed, this follows from [LaLu], Theorem 2.3.

To prove the proposition, we need the following lemma:

Lemma 6.3. *Let X and Y be varieties and E and F vector bundles on X and Y respectively. Then*

$$H^0(X \times Y, p^*E \otimes q^*F) = H^0(X, E) \otimes H^0(Y, F).$$

Proof. By the projection formula $p_*(p^*E \otimes q^*F) = E \otimes p_*q^*F$, where p_*q^*F is the trivial bundle on X with fiber $H^0(Y, F)$. Hence

$$\begin{aligned} H^0(X \times Y, p^*E \otimes q^*F) &= H^0(X, p_*(p^*E \otimes q^*F)) \\ &= H^0(X, E \otimes p_*q^*F) = H^0(X, E) \otimes H^0(Y, F). \end{aligned}$$

□

Proof of proposition.

i) Since Ψ^n is a polynomial in operations λ^j , it suffices to prove that $H^0(X, (\Omega_X^i)^{\otimes n}) = H^0(\tilde{X}, (\Omega_{\tilde{X}}^i)^{\otimes n})$. But this is well known (the proof is the same as that of [Ha] II 8.19).

ii) Consider the projections $X \xleftarrow{p} X \times Y \xrightarrow{q} Y$. We have

$$\Omega_{X \times Y}^1 = p^*\Omega_X^1 \oplus q^*\Omega_Y^1.$$

Hence

$$\Omega_{X \times Y}^m = \bigoplus_{i+j=m} p^*\Omega_X^i \otimes q^*\Omega_Y^j.$$

As $\bar{K}(X)$ is a special λ -ring,

$$\begin{aligned} \Psi^n \Omega_{X \times Y}^m &= \bigoplus_{i+j=m} \Psi^n p^*\Omega_X^i \otimes \Psi^n q^*\Omega_Y^j \\ &= \bigoplus_{i+j=m} p^*\Psi^n \Omega_X^i \otimes q^*\Psi^n \Omega_Y^j. \end{aligned}$$

It follows that

$$H^0(X \times Y, \Psi^n \Omega_{X \times Y}^m) = \bigoplus_{i+j=m} H^0(X, \Psi^n \Omega_X^i) \otimes H^0(Y, \Psi^n \Omega_Y^j),$$

i.e. $\mu_n(X \times Y) = \mu_n(X)\mu_n(Y)$.

iii) Since $\mathbb{P}^k$ is birational to $(\mathbb{P}^1)^k$, by i), ii) above it suffices to prove that $\mu_n(\mathbb{P}^1) = 1$. We have

$$\Psi^n \Omega_{\mathbb{P}^1}^i = \begin{cases} \mathcal{O}_{\mathbb{P}^1}, & \text{if } i = 0, \\ \mathcal{O}_{\mathbb{P}^1}(-2n), & \text{if } i = 1, \\ 0, & \text{if } i > 1. \end{cases}$$

and therefore $\mu_n(\mathbb{P}^1) = 1 \in M$. This proves the proposition. $\square$

Proposition 6.4. *Let X be a variety of dimension ≤ 2 . Then*

$$\mu(\text{Sym}^m X) = \lambda^m(\mu(X)).$$

Proof. The class of X in $K_0[\mathcal{V}_{\mathbb{C}}]$ is a linear combination of classes of smooth projective varieties of dimension 0, 1 and 2. Thus we may assume that X is smooth projective. If $\dim X = 0$ the assertion is trivial. We will prove the assertion in case $\dim X = 2$ (the case of curves is similar and simpler since $\text{Sym}^n X$ is smooth if $\dim X = 1$). So let X be a smooth projective surface.

We will use the smooth variety $\text{Hilb}^m X$ —the Hilbert scheme of zero-dimensional subschemes of X . It follows immediately from the main theorem of L. Goettsche [Go] that the classes of $\text{Hilb}^m X$ and $\text{Sym}^m X$ are equal in $K_0[\mathcal{V}_{\mathbb{C}}]/[\mathbb{A}^1]$. In particular,

$$\mu_n(\text{Hilb}^m X) = \mu_n(\text{Sym}^m X).$$

By Lemma 7.4 below $H^0(\text{Hilb}^m X, \Omega_{\text{Hilb}^m X}^i) = H^0(X^m, \Omega_{X^m}^i)^{S_m}$. Thus we need to prove the following equality in M :

$$\begin{aligned} 1 + H^0(X^m, \Omega^1)^{S_m} s + \cdots + H^0(X^m, \Omega^{2m})^{S_m} s^{2m} \\ = \lambda^m(1 + H^0(X, \Omega^1) s + H^0(X, \Omega^2) s^2). \end{aligned}$$

Recall from Example 4.2 that

$$\lambda^j(V s^i) = \begin{cases} \text{Sym}^j(V) s^{ij}, & \text{if } i \text{ is even,} \\ \Lambda^j(V) s^{ij}, & \text{if } i \text{ is odd.} \end{cases}$$

Let $\sum_i V_i s^i \in M$ be a graded vector space. Then

$$\lambda^m\left(\sum_i V_i s^i\right) = \sum_i \sum_{j_1 n_1 + \cdots + j_k n_k = i} \lambda^{n_1}(V_{j_1}) \otimes \cdots \otimes \lambda^{n_k}(V_{j_k}) s^i.$$

Let us prove the following general lemma. Let Y be a smooth projective variety and F a vector bundle on Y of rank r . Fix $m \geq 1$ and denote by $p_i : Y^m \rightarrow Y$ the projection to the i th factor. Put $F_i := p_i^* F$, $\mathcal{F} := F_1 \oplus \cdots \oplus F_m$.

Lemma 6.5. *There is a natural isomorphism of graded vector spaces*

$$\sum_j H^0(Y^m, \Lambda^j(\mathcal{F}))^{S_m} s^j = \lambda^m\left(\sum_j H^0(Y, \Lambda^j F) s^j\right).$$

Proof. We have

$$\Lambda^\bullet(\mathcal{F}) = \bigoplus_{(j_1, \dots, j_m)} \Lambda^{j_1} F_1 \otimes \dots \otimes \Lambda^{j_m} F_m,$$

By Lemma 6.3,

$$H^0(Y^m, \Lambda^{j_1} F_1 \otimes \dots \otimes \Lambda^{j_m} F_m) \simeq \bigotimes_{i=1}^m H^0(Y^m, \Lambda^{j_i} F_i) \simeq \bigotimes_{i=1}^m H^0(Y, \Lambda^{j_i} F).$$

Put $\mathcal{F}_{j_1 \dots j_m} = \Lambda^{j_1} F_1 \otimes \dots \otimes \Lambda^{j_m} F_m$. Consider the S_m -action on the sheaf $\Lambda^\bullet(\mathcal{F})$. The sheaves $\mathcal{F}_{j_1 \dots j_m}$ and $\mathcal{F}_{j'_1 \dots j'_m}$ are in the same S_m -orbit if and only if the multisets $\{j_1, \dots, j_m\}$ and $\{j'_1, \dots, j'_m\}$ are equal. For each multiset $\{j_1, \dots, j_m\}$ choose a representative $(j_1, \dots, j_m)$ of the corresponding S_m -orbit. Put $G_{j_1 \dots j_m} = \text{Stab}_{S_m}(j_1, \dots, j_m)$. Then

$$W = H^0(Y^m, \Lambda^\bullet \mathcal{F})^{S_m} = \sum_{\{j_1, \dots, j_m\}} H^0(Y^m, \mathcal{F}_{j_1 \dots j_m})^{G_{j_1 \dots j_m}}.$$

Consider the space

$$H^0(Y^m, \mathcal{F}_{j_1 \dots j_m})^{G_{j_1 \dots j_m}} = (\otimes_s H^0(Y^m, \Lambda^{j_s} F_s))^{G_{j_1 \dots j_m}}.$$

Assume for simplicity of notation that the multiset $\{j_1, \dots, j_m\}$ contains k different elements:

$$j_1 = \dots = j_{t_1} \neq j_{t_1+1} = \dots = j_{t_1+t_2} \neq \dots = j_{t_1+\dots+t_k}.$$

Then $G_{j_1 \dots j_m} = S_{t_1} \times \dots \times S_{t_k}$. For example, the subgroup S_{t_1} acts by permuting factors in $\Lambda^{j_1} F_1 \otimes \dots \otimes \Lambda^{j_1} F_{t_1}$. A transposition of two factors in this tensor product corresponds to a product of j_1 transpositions on the level of $\mathcal{F}^{\otimes j_1 t_1}$. Therefore,

$$H^0(Y^m, \Lambda^{j_1} F_1 \otimes \dots \otimes \Lambda^{j_{t_1}} F_{t_1})^{S_{t_1}} \simeq \begin{cases} \text{Sym}^{t_1} H^0(Y, \Lambda^{j_1} F), & \text{if } j_1 \text{ is even,} \\ \Lambda^{t_1} H^0(Y, \Lambda^{j_1} F), & \text{if } j_1 \text{ is odd,} \end{cases}$$

The space $H^0(Y^m, \mathcal{F}_{j_1 \dots j_m})^{G_{j_1 \dots j_m}}$ is the tensor product of k factor of the form $\text{Sym}^{t_p} H^0(Y, \Lambda^{l_p} F)$ (if l_p is even) or $\Lambda^{t_p} H^0(Y, \Lambda^{l_p} F)$ (if l_p is odd). Also the degree of this space is equal to $\sum_s j_s$. This proves the lemma. $\square$

Now apply the lemma with $Y = X$ and $F = \Omega_X^1$ to get the proposition. $\square$

Remark 6.6. Unfortunately, the assertion of the last proposition is no longer true if we replace the measure μ by μ_n for $n \geq 2$. A counter example is provided by a smooth projective curve of genus ≥ 2 . Indeed, then $\text{Sym}^m X$ is stably birational to the Jacobian of X and hence $\mu_n(\text{Sym}^m X) = \mu(\text{Sym}^m X)$. On the other hand it is clear that $\mu_n(X)$ and hence $\lambda^m(\mu_n(X))$ depends on n .

Question 6.7. Is $\mu : K_0[\mathcal{V}_{\mathbb{C}}] \rightarrow \mathbb{Z}[M]$ is a λ -homomorphism? It seems likely to us that the answer is affirmative. We could prove it if we knew that for any smooth projective variety Z and any $m \geq 1$ the class of $\text{Sym}^m Z$ is equal in $K_0[\mathcal{V}_{\mathbb{C}}]/[\mathbb{A}^1]$ to the class of a resolution of $\text{Sym}^m Z$. So one might generalize and ask the following question. Let X be a non-singular complex projective variety with an action by a finite group G , and Y a non-singular projective variety birationally equivalent to X/G . Is it always true that

$$[X/G] \equiv [Y] \pmod{[\mathbb{A}^1]}?$$

7. IRRATIONALITY THEOREMS

Let X be a smooth projective surface. Let $m \geq 1$ be an integer and $\text{Hilb}^m X$ be the Hilbert scheme of zero-dimensional subschemes of X of length m . It is well known that the natural (Hilbert-Chow) morphism $g : \text{Hilb}^m X \rightarrow \text{Sym}^m X$ is a resolution of singularities which is a semismall map ([Na] 1.15, 6.10). It follows immediately from the main theorem of L. Gottsche [Go] that the classes of $\text{Hilb}^m X$ and $\text{Sym}^m X$ are equal in $K_0[\mathcal{V}_{\mathbb{C}}]/[\mathbb{A}^1]$. Thus for all $n \geq 1$

$$\mu_n(\text{Hilb}^m X) = \mu_n(\text{Sym}^m X).$$

Let us introduce some notation. Consider the pullback diagram

$$\begin{array}{ccc} Y & \xrightarrow{f} & X^m \\ \pi \downarrow & & \downarrow \pi \\ \text{Hilb}^m X & \xrightarrow{g} & \text{Sym}^m X, \end{array}$$

where the vertical arrows are quotient morphisms by the S_m -action. Let $\text{Sym}^m X_*$ denote the open subspace of $\text{Sym}^m X$ consisting of all multisets $\{x_1, \dots, x_m\}$ in which at least $m - 1$ points are distinct. Pulling back the previous diagram to $\text{Sym}^m X_*$, we have

$$\begin{array}{ccc} Y_* & \xrightarrow{f} & X_*^m \\ \pi \downarrow & & \downarrow \pi \\ \text{Hilb}^m X_* & \xrightarrow{g} & \text{Sym}^m X_*. \end{array}$$

Denote by Δ_* the intersection of the “fat” diagonal $\Delta \subset X^m$ with X_*^m . Then Y_* is the blowup of X_*^m along the smooth subvariety Δ_* which is of codimension 2. So Y_* is smooth. Let $E \subset Y_*$ be the exceptional divisor. The map $\pi : Y_* \rightarrow \text{Hilb}^m X_*$ is unramified away from E and has ramification of degree 2 along E . Since the map g is semismall, the complement of the open subset $\text{Hilb}^m X_* \subset \text{Hilb}^m X$ has codimension ≥ 2 . Hence also $Y \setminus Y_*$ has codimension ≥ 2 in Y .

Finally, let $\text{Sym}_+^m X$ be the image of $X^m - \Delta$. We pull back the previous diagram to $\text{Sym}_+^m X$:

$$(7.1) \quad \begin{array}{ccc} Y_+ & \xrightarrow{f} & X_+^m \\ \pi \downarrow & & \downarrow \pi \\ \text{Hilb}^m X_+ & \xrightarrow{g} & \text{Sym}^m X_+. \end{array}$$

The vertical maps π are étale and the horizontal maps f, g are isomorphisms. The complement of the open subset $X_+^m \subset X^m$ has codimension 2.

Lemma 7.1. *For any $m, n \geq 1$ we have*

$$H^0(\text{Hilb}^m X, (\Omega_{\text{Hilb}^m X}^1)^{\otimes n}) \subset H^0(X^m, (\Omega_{X^m}^1)^{\otimes n})^{S_m}.$$

Proof. Consider the diagram (7.1). We have

$$\begin{aligned} H^0(\text{Hilb}^m X, (\Omega_{\text{Hilb}^m X}^1)^{\otimes n}) &\subset H^0(\text{Hilb}^m X_+, (\Omega_{\text{Hilb}^m X_+}^1)^{\otimes n}) \\ &= H^0(Y_+, (\Omega_{Y_+}^1)^{\otimes n})^{S_m} = H^0(X_+^m, (\Omega_{X_+^m}^1)^{\otimes n})^{S_m}. \end{aligned}$$

Moreover,

$$H^0(X_+^m, (\Omega_{X_+^m}^1)^{\otimes n})^{S_m} = H^0(X^m, (\Omega_{X^m}^1)^{\otimes n})^{S_m},$$

since X_+^m has complement of codimension 2 in X^m . $\square$

Lemma 7.2. *For any $m, n \geq 1$ we have*

$$H^0(\text{Hilb}^m X, \omega_{\text{Hilb}^m X}^{\otimes n}) = H^0(X^m, \omega_{X^m}^{\otimes n})^{S_m}.$$

Proof. It suffices to prove that

$$H^0(\text{Hilb}^m X_*, \omega_{\text{Hilb}^m X_*}^{\otimes n}) = H^0(X_*^m, \omega_{X_*^m}^{\otimes n})^{S_m}.$$

We have the natural injective maps

$$\alpha : \pi^* \omega_{\text{Hilb}^m X_*} \rightarrow \omega_{Y_*}, \quad \beta : f^* \omega_{X_*^m} \rightarrow \omega_{Y_*}.$$

It suffices to prove that $\text{im}(\alpha) = \text{im}(\beta)$. Indeed,

$$H^0(Y_*, \pi^* \omega_{\text{Hilb}^m X_*})^{S_m} = H^0(\text{Hilb}^m X_*, \omega_{\text{Hilb}^m X_*}).$$

Both α and β are surjective away from E . So it remains to analyze the maps g and π near E .

Choose a point $p = (a_1, a_2 = a_1, a_3, \dots, a_m) \in \Delta_*$ and $q \in g^{-1}(p)$. There exist local (analytic) coordinates $x_1, x_2, \dots, x_{2m}$ near p and local coordinates $y_1, \dots, y_{2m}$ near q so that

$$g^*(x_1) = y_1, \quad g^*(x_2) = y_1 y_2, \quad g^*(x_3) = y_3, \quad \dots \quad g^*(x_{2m}) = y_{2m}.$$

Then $y_1 = 0$ is the local equation of E . Thus

$$g^*(dx_1 \wedge \dots \wedge dx_{2m}) = y_1 dy_1 \wedge \dots \wedge y_{2m}$$

and

$$\beta((dx_1 \wedge \cdots \wedge dx_{2m})^{\otimes n}) = y_1^n (dy_1 \wedge \cdots \wedge y_{2m})^{\otimes n},$$

so that $\text{im}(\beta) = \omega_{Y_*}^{\otimes n}(-nE)$. Similarly, we can define local coordinates $z_1, \dots, z_{2m}$ near $\pi(q) \in \text{Hilb}^m X_*$, so that

$$\pi^*(z_1) = y_1^2, \pi^*(z_2) = y_2, \dots, \pi^*(z_{2m}) = y_{2m}.$$

Therefore

$$g^*(dz_1 \wedge \cdots \wedge dz_{2m}) = 2y_1 dy_1 \wedge \cdots \wedge y_{2m}$$

and

$$\alpha((dz_1 \wedge \cdots \wedge dz_{2m})^{\otimes n}) = 2^n y_1^n (dy_1 \wedge \cdots \wedge y_{2m})^{\otimes n}.$$

That means $\text{im}(\alpha) = \omega_{Y_*}(-nE)$. □

Lemma 7.3. *For any $m, n \geq 1$ we have*

$$H^0(X^m, \omega_{X^m}^{\otimes n})^{S_m} = \text{Sym}^m H^0(X, \omega_X^{\otimes n}).$$

Proof. Let $p_i : X^m \rightarrow X$ denote the projection on the i th factor. Then

$$\omega_{X^m}^{\otimes n} = (p_1^* \omega_X \otimes \cdots \otimes p_m^* \omega_X)^{\otimes n} \simeq p_1^* \omega_X^{\otimes n} \otimes \cdots \otimes p_m^* \omega_X^{\otimes n},$$

and hence

$$H^0(X^m, \omega_{X^m}^{\otimes n}) = H^0(X, \omega_X^{\otimes n})^{\otimes m}.$$

The S_m -action permutes the factors and

$$H^0(X^m, \omega_{X^m}^{\otimes n})^{S_m} = \text{Sym}^m H^0(X, \omega_X^{\otimes n}),$$

since $\dim X$ is even. □

Lemma 7.4. *For any $m, i \geq 1$ we have*

$$H^0(\text{Hilb}^m X, \Omega_{\text{Hilb}^m X}^i) = H^0(X^m, \Omega_{X^m}^i)^{S_m}.$$

Proof. Since Y_* is a blowup of X_*^m along a smooth subvariety we have $H^0(Y_*, \Omega_{Y_*}^i) = H^0(X_*^m, \Omega_{X_*^m}^i)$. Also $H^0(X_*^m, \Omega_{X_*^m}^i) = H^0(X^m, \Omega_{X^m}^i)$, since X_*^m has complement of codimension ≥ 2 in X^m . Thus we must show that

$$H^0(\text{Hilb}^m X, \Omega_{\text{Hilb}^m X}^i) = H^0(Y_*, \Omega_{Y_*}^i)^{S_m}.$$

Let $q \in Y_*$, $G_q = \text{Stab}(q) \subset S_m$. It suffices to show that

$$(\pi^{-1} \Omega_{\text{Hilb}^m X_*})_q = (\Omega_{Y_*, q})^{G_q}.$$

If $q \in Y_+$, then $G_q = 1$ and $(\pi^{-1} \Omega_{\text{Hilb}^m X_*})_q = (\Omega_{Y_*, q})$. Suppose $q \in E$ and hence $G_q = \mathbb{Z}/2\mathbb{Z}$. Consider local coordinates $y_1, \dots, y_{2m}$ at q and local coordinates $z_1, \dots, z_{2m}$ at $\pi(q)$ as in the proof of Lemma 7.2 above, so that

$$\pi^*(z_1) = y_1^2, \pi^*(z_2) = y_2, \dots, \pi^*(z_{2m}) = y_{2m}.$$

An element $y_1^k dy_1$ is G_q -invariant if and only if k is odd. The lemma follows. □

Proposition 7.5. *Fix integers $n, i \geq 1$. Then the absolute value of the integer $h_n^i(\text{Hilb}^m X)$ is bounded independently of m .*

Proof. Recall that $h_n^i(Z) = H^0(Z, \Psi^n \Omega_Z^i)$. The Adams operation Ψ^n is a polynomial in the operations λ^i . Hence it suffices to prove that for all $n \geq 0$, the dimension of the space $H^0(\text{Hilb}^m X, (\Omega_{\text{Hilb}^m X}^1)^{\otimes n})$ is bounded independently of m . By Lemma 7.1 above

$$H^0(\text{Hilb}^m X, (\Omega_{\text{Hilb}^m X}^1)^{\otimes n}) \subset H^0(X^m, (\Omega_{X^m}^1)^{\otimes n})^{S_m}.$$

We will prove that $\dim H^0(X^m, (\Omega_{X^m}^1)^{\otimes n})^{S_m}$ is bounded independently of m .

Let $p_i : X^m \rightarrow X$ be the projection to the i th factor. Denote $F_i = p_i^* \Omega_X^1$. Then $\Omega_{X^m}^1 = \bigoplus_{i=1}^m F_i$ and

$$(\Omega_{X^m}^1)^{\otimes n} = \bigoplus_{i_1, \dots, i_n \in \{1, \dots, m\}} F_{i_1} \otimes \dots \otimes F_{i_n}.$$

The S_m -action permutes the summands $F_{i_1} \otimes \dots \otimes F_{i_n}$ and the orbits correspond to partitions P of the set $\{1, \dots, n\}$. Thus

$$H^0(X^m, \Omega^{\otimes n})^{S_m} = \bigoplus_P H^0(X^m, F_P)^{\text{Stab}(P)},$$

Where F_P is one of the summands in the orbit, corresponding to P . Fix a partition P . Assume for simplicity of notation that P divides $\{1, \dots, n\}$ into k segments $P_1, \dots, P_k$, where $a < b$ for each $a \in P_s, b \in P_t$ if $s < t$. Let $\alpha_i = |P_i|$. Then

$$H^0(X^m, F_P) = H^0(X, \Omega^{\otimes \alpha_1}) \otimes \dots \otimes H^0(X, \Omega^{\otimes \alpha_k}).$$

Therefore $\dim H^0(X^m, F_P)$ is bounded independently of m . $\square$

Theorem 7.6. *Let X be a smooth projective surface of Kodaira dimension ≥ 0 . Then the zeta-function $\zeta_X(t)$ is not pointwise rational.*

Proof. Let F denote the field of fractions of $\mathbb{Z}[M]$. We will show that there exists $n \geq 1$ such that the power series

$$1 + \sum_{m=1}^{\infty} \mu_n(\text{Sym}^m X) t^m \in F[[t]]$$

is not rational.

First we claim that there exists $n \geq 1$ such that

$$h_n^{2d}(\text{Hilb}^m X) = \dim H^0(\text{Hilb}^m X, \omega_{\text{Hilb}^m X}^{\otimes n}) > 0$$

for all $m \geq 1$. Indeed, by our assumption on X there exists $n \geq 1$, such $H^0(X, \omega^{\otimes n}) \neq 0$ and by Lemmas 7.2 and 7.3,

$$H^0(\text{Hilb}^m X, \omega_{\text{Hilb}^m X}^{\otimes n}) = \text{Sym}^m H^0(X, \omega_X^{\otimes n})$$

for all $m \geq 1$. Fix one such n and consider the motivic measure $\mu_n : K_0[\mathcal{V}_{\mathbb{C}}] \rightarrow \mathbb{Z}[M] \subset F$. We have $\mu_n(\text{Sym}^m X) = \mu_n(\text{Hilb}^m X)$. It follows that

$\mu_n(\mathrm{Sym}^m X) \in M$ is a polynomial with constant term 1 and leading term $h_n^{2d}(\mathrm{Hilb}^m X)s^{2d}$.

Consider the group completion G of the monoid M . By Lemma 4.5, G is a free abelian group. Hence we may apply Proposition 2.6 to the power series $1 + \sum_{m=1}^{\infty} \mu_n(\mathrm{Sym}^m X)t^m$. Denote $g_m = \mu_n(\mathrm{Sym}^m X) \in M$. Assume that this power series is rational. Then there exist $k, i \geq 1$ and an element $g \in G$ such that

$$g_{i+(\alpha+1)k} = gg_{i+\alpha k}$$

for all $\alpha \geq 0$. The element g is a rational function in s . Since all coefficients g_m are nonzero polynomials in s it follows that g is also a nonzero polynomial. Note that g is not a monomial since the degrees of the polynomials g_m grow and their constant term is 1. But then the coefficients of fixed powers of s in the g_m cannot stay bounded, which contradicts Proposition 7.5. $\square$

8. THE SPECIAL GROTHENDIECK λ -RING OF VARIETIES.

As pointed out in Example 4.6, the symmetric power operations Sym^n define a λ -ring structure on the Grothendieck ring $K_0[\mathcal{V}_{\mathbb{C}}]$. The zeta-function $\zeta_X(t)$ looks formally like a universal λ -homomorphism, but in fact it is not a ring homomorphism at all. To see this, it suffices to note that $\zeta_X(t)$ is a ratio of polynomials with constant term 1 for every curve X ; thus (4.1) implies that $\zeta_X(t) \cdot \zeta_Y(t)$ is rational, while Theorem 7.6 implies $\zeta_{X \times Y}(t)$ is irrational whenever X and Y both have positive genus.

Let $\{\lambda^n\}$ denote the λ -structure opposite to $\{\mathrm{Sym}^n\}$. This seems to be the natural choice of λ -structure on $K_0[\mathcal{V}_{\mathbb{C}}]$ insofar as $X \mapsto \mathrm{Sym}^n X$ behaves like a symmetric power map, for instance on cohomology. The choice does not affect which classes are virtually finite, but it will make a difference when we specialize, since, as we have seen, specialization does not commute with taking opposites.

Lemma 8.1. *If X is a variety over $\mathbb{C}$ whose image in $K_0[\mathcal{V}_{\mathbb{C}}]$ is virtually finite, then $\zeta_X(t)$ is globally rational.*

Proof. Let $[X] = y - z$, where y and z are finite. As λ_t is a ring homomorphism,

$$\zeta_X(T) = \lambda_{-t}([X])^{-1} = \lambda_{-t}(y)^{-1} \lambda_{-t}(z)$$

is globally rational in $K_0[\mathcal{V}_{\mathbb{C}}]$. $\square$

We have the following variant of Theorem 3.7:

Proposition 8.2. *The class of any 1-dimensional variety in $K_0[\mathcal{V}_{\mathbb{C}}]$ is virtually finite.*

Proof. If Y is a closed subvariety of X with complement U , then if any two of X , Y , and U are virtually finite, the third is so as well. A point Z is finite dimensional (in fact 1-dimensional in the λ -ring sense). Thus,

we reduce exactly as in the proof of Theorem 3.7 to the case of a single projective non-singular curve X . Now,

$$\lambda_t([\mathbb{P}^1]) = (1+t)(1 + [\mathbb{A}^1]t),$$

so $\mathbb{P}^1$ is 2-dimensional. Writing $X = \mathbb{P}^1 - (\mathbb{P}^1 - X)$, (3.2) implies

$$\lambda_t(\mathbb{P}^1 - X) = (1+t)(1 + \mathbb{A}^1 t) \lambda_t(X)^{-1} = (1+t)(1 + \mathbb{A}^1 t) \zeta_X(-t) \in K_0[\mathcal{V}_K][t].$$

The proposition follows. $\square$

Corollary 8.3. *The virtual finiteness of a complex surface X depends only on the birational class of X .*

Definition 8.4. We call the specialization of the Grothendieck ring $K_0[\mathcal{V}_K]$ with respect to λ^n the **Grothendieck λ -ring** of K and denote it $K_0[\mathcal{V}_K]_\sigma$. The image of $\zeta_X(t)$ in $K_0[\mathcal{V}_K]_\sigma$ will be denoted $\zeta_{X\sigma}(t)$

In any special λ -ring, the set of virtually finite elements is clearly a λ -subring. In particular, we have:

Proposition 8.5. *For every $X/\mathbb{C}$ which is virtually finite in $K_0[\mathcal{V}_\mathbb{C}]$ and every positive integer n , $\text{Sym}^n X$ is again virtually finite in $K_0[\mathcal{V}_\mathbb{C}]$.*

Proof. The identities relating symmetric and exterior powers show that $\text{Sym}^n X$ lies in the λ -subring of $K_0[\mathcal{V}_\mathbb{C}]_\sigma$ generated by X . $\square$

Proposition 8.6. *Every principally polarized abelian surface $X/\mathbb{C}$ is virtually finite in $K_0[\mathcal{V}_\mathbb{C}]_\sigma$.*

Proof. It is well known that every principally polarized abelian variety of dimension ≤ 3 is a product of Jacobian varieties. (We do not know who first made this observation, but it is an immediate consequence of Torelli's theorem and Baily's theorem [Ba].) It suffices, therefore, to prove that the Jacobian J of a genus 2 curve X is virtually finite. The map $\text{Sym}^2 X \rightarrow J$ is a birational equivalence, so the proposition follows from Corollary 8.3. $\square$

We remark that the fact that sufficiently high symmetric powers of a non-singular projective curve are projective space bundles over its Jacobian variety does not immediately imply virtual finiteness of Jacobians, since it is not obvious that the virtual finiteness of a projective space bundle or even a vector bundle over a given variety implies the virtual finiteness of that variety. However, it is easy to prove the pointwise rationality of $\zeta_{J\sigma}(t)$ for Jacobians J .

Question 8.7. Is $\zeta_{A\sigma}(t)$ rational (globally or pointwise) for all abelian varieties A ?

More optimistically, we might ask:

Question 8.8. Is $\zeta_{X\sigma}(t)$ rational for all varieties X ? Is $K_0[\mathcal{V}_\mathbb{C}]_\sigma$ finite dimensional in the sense of Definition 4.7?

REFERENCES

- [AT] M. Atiyah, D. Tall: Group representations, λ -rings, and the J -homomorphism, *Topology* **8** (1969), 253–297.
- [Ba] W. Baily: On the moduli of Jacobian varieties, *Annals of Math.* **71** (1960), 303–314.
- [Bi] F. Bittner: The universal Euler characteristic for varieties of characteristic zero, arXiv: math.AG/0111062.
- [Bl] S. Bloch: K^2 of Artinian $\mathbb{Q}$ -algebras, with application to algebraic cycles. *Comm. Algebra* **3** (1975), 405–428.
- [BKL] S. Bloch, A. Kas, D. Lieberman: Zero cycles on surfaces with $p_g = 0$. *Compositio Math.* **33** (1976), 135–145.
- [A.B] A. Borel: *Linear Algebraic Groups*, 2nd ed., Springer-Verlag, New York, 1991.
- [E.B] E. Borel: Sur une application d’un théorème de M. Hadamard, *Bull. Sci. Mathématiques*, Ser. 2 **18** (1894), 22–25.
- [FL] W. Fulton, S. Lang: *Riemann-Roch Algebra*, Springer-Verlag, 1985.
- [Go] L. Goettsche: On the motive of the Hilbert scheme of points on a complex surface. *Math. Res. Lett.* **8** (2001), 613–627.
- [Ha] R. Hartshorne: *Algebraic Geometry*, Springer-Verlag, 1977.
- [Ka] M. Kapranov: The elliptic curve in the S-duality theory and Eisenstein series for Kac-Moody groups, arXiv: math.AG/0001005.
- [LL] M. Larsen and V. Lunts: Motivic measures and stable birational geometry, arXiv: math.AG/0110255, to appear in *Mosc. Math. J.*
- [Mu] D. Mumford: Rational equivalence of 0-cycles on surfaces. *J. Math. Kyoto Univ.* **9** (1968), 195–204.
- [Na] H. Nakajima: *Lectures on Hilbert Schemes of Points on Surfaces*, AMS Univ. Lect. Ser. Vol 18, 1999.
- [Po] B. Poonen: The Grothendieck ring of varieties is not a domain, *Math. Res. Lett.* **9** (2002), 493–498.
- [Re] M. Reid: Chapters on Algebraic Surfaces, in *Complex Algebraic Geometry*, ed. J. Kollár, American Mathematical Society, Providence, 1997.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405, USA

E-mail address: larsen@math.indiana.edu

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405, USA

E-mail address: vlunts@indiana.edu